

## What families can do to help their teens achieve a healthy weight and positive body image:\*

Cornerstone Behavior	Specific Steps that Parents/Families Can Take	Supportive Research
Model healthy behaviors for your children	<ul style="list-style-type: none"> <li>• Avoid dieting or, at least, unhealthy dieting behaviors</li> <li>• Avoid making weight-related comments as much as possible</li> <li>• Engage in regular physical activity that you enjoy</li> <li>• Model healthy (but not perfect) eating patterns and food choices</li> </ul>	<ul style="list-style-type: none"> <li>• Although not consistent across studies, correlations are often seen between parental and adolescent eating, activity, and dieting behaviors. Associations are strongest when behaviors are noticed by teens (i.e., reported by teens rather than by parents) <sup>1-3</sup>.</li> <li>• Parents' comments about weight may be more important than their behaviors <sup>4-6</sup>. In the AHEAD study, negative weight comments within families were one of the strongest correlates of unhealthy weight control practices in teens with type 1 diabetes <sup>7</sup>.</li> </ul>
Provide an environment that makes it easy for your children to make healthy choices.	<ul style="list-style-type: none"> <li>• Make healthy foods readily available</li> <li>• Establish family meal norms that work for your family</li> <li>• Make physical activity the norm in your family, and limit TV watching</li> <li>• Support your teen's efforts to get involved in physical activity</li> </ul>	<ul style="list-style-type: none"> <li>• In Project EAT, the strongest correlate of fruit/vegetable intake was home availability <sup>8</sup></li> <li>• In focus groups, the teens stated very clearly: "If it's out, I'll eat it!" <sup>9</sup></li> <li>• Family meals are strongly correlated with better dietary quality <sup>10, 11</sup></li> <li>• Research suggests the importance of decreasing sedentary activity and increasing physical activity for weight management and the important role that parents can play in supporting their children's efforts to be active <sup>12-14</sup></li> </ul>
Focus less on weight; instead, focus on behaviors and overall health.	<ul style="list-style-type: none"> <li>• Encourage your teen to adopt healthy behaviors without focusing on weight loss</li> <li>• Help your teen develop an identity that goes beyond physical appearance</li> <li>• Establish a no-tolerance policy for weight teasing in your home</li> </ul>	<ul style="list-style-type: none"> <li>• Dieting for weight loss is associated with weight gain over time and the onset of disordered eating/eating disorders, suggesting the importance of focusing on more effective behaviors that can be implemented safely over a long period of time <sup>15-19</sup></li> <li>• Among teens, body image and more global self-esteem tend to be linked and there is a high prevalence of body dissatisfaction among teens <sup>20-22</sup></li> </ul>
Provide a supportive environment with lots of talking and even more listening	<ul style="list-style-type: none"> <li>• Be there to listen and provide support when your teen discusses weight concerns</li> <li>• When your teen talks about fat, find out what's really going on</li> <li>• Keep the communication lines open – no matter what</li> <li>• Provide unconditional love, not love based on weight, and let your child know how you feel</li> </ul>	<ul style="list-style-type: none"> <li>• The prevalence of weight-teasing is high among teens, particularly overweight teens <sup>23, 24</sup></li> <li>• The impact of teasing on psychological well being and disordered eating was found to be high in Project EAT <sup>23, 25</sup></li> <li>• High levels of family communication protect adolescents against an array of health-compromising behaviors <sup>7, 26</sup></li> <li>• In individual interviews, overweight teens indicated that in the face of weight-related teasing/mistreatment, they appreciated support from their family, not weight-loss advice <sup>24, 27</sup></li> </ul>

\*Source:

Neumark-Sztainer D. "I'm, Like, SO Fat!" Helping Your Teen Make Healthy Choices about Eating and Exercise in a Weight-Obsessed World. New York: The Guilford Press; 2005. Reprinted with permission of the Guilford Press. As reproduced in: Neumark-Sztainer D. Preventing the broad spectrum of weight-related problems: working with parents to help teens achieve a healthy weight and a positive body image. *Journal of Nutrition Education & Behavior*. 2005; 37 (Suppl 2):S133-40.

## References

1. Benedikt R, Wertheim E, Love A. Eating attitudes and weight-loss attempts in female adolescents and their mothers. *Journal of Youth and Adolescence*. 1998;27(1):43-57.
2. Hanson NI, Neumark-Sztainer D, Eisenberg ME, Story M, Wall M. Associations between parental report of the home food environment and adolescent intakes of fruits, vegetables, and dairy foods. *Public Health Nutr*. 2005;8:77-85.
3. Sallis JF, Prochaska JJ, Taylor WC. A review of correlates of physical activity of children and adolescents. *Med Sci Sports Exerc*. 2000;32(5):963-975.
4. Wertheim EH, Mee V, Paxton SJ. Relationships among adolescent girls' eating behaviors and their parents' weight-related attitudes and behaviors. *Sex Roles*. 1999;41:169-187.
5. Fulkerson JA, McGuire MT, Neumark-Sztainer D, Story M, French SA, Perry CL. Weight-related attitudes and behaviors of adolescent boys and girls who are encouraged to diet by their mothers. *Int J Obes*. 2002;26:1579-1587.
6. Dixon R, Adair V, O'Connor S. Parental influences of the dieting beliefs and behaviors of adolescent females in New Zealand. *J Adolesc Health*. 1996;19:303-307.
7. Mellin AE, Neumark-Sztainer D, Patterson J, Sockalosky J. Unhealthy weight management behavior among adolescent girls with type 1 diabetes mellitus: The role of familial eating patterns and weight-related concerns. *J Adolesc Health*. 2004;35(4):278-289.
8. Neumark-Sztainer D, Wall MM, Hannan PJ, Story M, Croll J, Perry C. Correlates of fruit and vegetable intake among adolescents: Findings from Project EAT. *Prev Med*. 2003;37(3):198-208.
9. Neumark-Sztainer D, Story M, Perry C, Casey MA. Factors influencing food choices of adolescents: Findings from focus-group discussions with adolescents. *J Am Diet Assoc*. 1999;99(8):929-937.
10. Neumark-Sztainer D, Hannan PJ, Story M, Croll J, Perry C. Family meal patterns: Associations with sociodemographic characteristics and improved dietary intake among adolescents. *J Am Diet Assoc*. 2003;103(3):317-322.
11. Gillman MW, Rifas-Shiman SL, Frazier AL, et al. Family dinner and diet quality among older children and adolescents. *Arch Fam Med*. 2000;9(3):235-240.
12. Utter J, Neumark-Sztainer D, Jeffery R, Story M. Couch potatoes or french fries: Are sedentary behaviors associated with body mass index, physical activity, and dietary behaviors among adolescents? *J Am Diet Assoc*. 2003;103(10):1298-1305.
13. McGuire M, Hannan P, Neumark-Sztainer D, Falkner-Cossrow N, Story M. Parental correlates of physical activity in a racially/ethnically-diverse adolescent sample. *J Adolesc Health*. 2002;30(4):253-261.
14. Wing RR, Hill JO. Successful weight loss maintenance. *Annual Reviews of Nutrition*. 2001;21:323-341.
15. Neumark-Sztainer D, Wall M, Guo J, Story M, Haines J, Eisenberg M. Obesity, disordered eating, and eating disorders in a longitudinal study of adolescents: How do dieters fare five years later? *Journal of the American Dietetic Association*. Under Review.
16. Field AE, Austin SB, Taylor CB, et al. Relation between dieting and weight change among preadolescents and adolescents. *Pediatrics*. 2003;112(4):900-906.
17. Stice E, Cameron RP, Killen JD, Hayward C, Taylor CB. Naturalistic weight-reduction efforts prospectively predict growth in relative weight and onset of obesity among female adolescents. *J Consult Clin Psychol*. 1999;67(6):967-974.
18. Patton GC, Selzer R, Coffey C, Carlin JB, Wolfe R. Onset of adolescent eating disorders: Population based cohort study over 3 years. *BMJ*. Mar 20 1999;318(7186):765-768.
19. Stice E, Presnell K, Shaw H, Rohde P. Psychological and behavioral risk factors for obesity onset in adolescent girls: A prospective study. *J Consult Clin Psychol*. 2005;73(2):195-202.

20. Ricciardelli LA, McCabe MP. Psychometric evaluation of the Body Change Inventory: An assessment instrument for adolescent boys and girls. *Eat Behav.* Spring 2002;3(1):45-59.
21. Wertheim E, Koerner J, Paxton S. Longitudinal predictors of restrictive eating and bulimic tendencies in three different age groups of adolescent girls. *Journal of Youth and Adolescence.* 2001;30(1):69-81.
22. Wichstrom L. The emergence of gender difference in depressed mood during adolescence: The role of intensified gender socialization. *Dev Psychol.* 1999;35(1):232-245.
23. Neumark-Sztainer D, Falkner N, Story M, Perry C, Hannan PJ, Mulert S. Weight-teasing among adolescents: Correlations with weight status and disordered eating behaviors. *Int J Obes Relat Metab Disord.* 2002;26(1):123-131.
24. Neumark-Sztainer D, Story M, Faibisch L. Perceived stigmatization among overweight African American and Caucasian adolescent girls. *J Adolesc Health.* 1998;23(5):264-270.
25. Eisenberg ME, Neumark-Sztainer D, Story M. Associations of weight-based teasing and emotional well-being among adolescents. *Arch Pediatr Adolesc Med.* 2003;157:733-738.
26. Resnick MD, Bearman PS, Blum RW, et al. Protecting adolescents from harm: Findings from the National Longitudinal Study on Adolescent Health. *JAMA.* 1997;278(10):823-832.
27. Neumark-Sztainer D. *"I'm, Like, SO Fat!": Helping Your Teen Make Healthy Choices about Eating and Exercise in a Weight-Obsessed World.* New York: The Guilford Press; 2005.